

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use SPRIX® safely and effectively. See full prescribing information for SPRIX®.

SPRIX® (ketorolac tromethamine) Nasal Spray
Initial U.S. Approval: 1989

WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS

See full prescribing information for complete boxed warning.

- **Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in treatment and may increase with duration of use (5.1)**
- **SPRIX® is contraindicated in the setting of coronary artery bypass graft (CABG) surgery (4, 5.1)**
- **NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious GI events (5.2)**

RECENT MAJOR CHANGES

Indications and Usage (1) 01/2018

INDICATIONS AND USAGE

SPRIX is a nonsteroidal anti-inflammatory drug indicated in adult patients for the short term (up to 5 days) management of moderate to moderately severe pain that requires analgesia at the opioid level. (1)

DOSAGE AND ADMINISTRATION

- Use the lowest effective dosage for shortest duration consistent with individual patient treatment goals. (2.1)
- SPRIX is not an inhaled product. For adult patients < 65 years of age: 31.5 mg (one 15.75 mg spray in each nostril) every 6 to 8 hours. The maximum daily dose is 126 mg. (2.2, 2.3)
- For patients ≥ 65 years of age, renally impaired patients, and patients less than 50 kg (110 lbs): 15.75 mg (one 15.75 mg spray in only one nostril) every 6 to 8 hours. The maximum daily dose is 63 mg. (2.4)
- SPRIX nasal spray should be discarded within 24 hours of taking the first dose, even if the bottle still contains some medication. (2.5)

DOSAGE FORMS AND STRENGTHS

SPRIX (ketorolac tromethamine) Nasal Spray: 15.75 mg of ketorolac tromethamine in each 100 µL spray. Each 1.7 g bottle contains 8 sprays. (3)

CONTRAINDICATIONS

- Known hypersensitivity to ketorolac or any components of the drug product (4)
- History of asthma, urticaria, or other allergic-type reactions after taking aspirin or other NSAIDs (4)
- In the setting of CABG surgery (4)
- Use in patients with active peptic ulcer disease or with recent GI bleeding or perforation (4)
- Use as a prophylactic analgesic before any major surgery (4)
- Use in patients with advanced renal disease or patients at risk for renal failure due to volume depletion (4)
- Use in patients with suspected or confirmed cerebrovascular bleeding, patients with hemorrhagic diathesis, incomplete hemostasis, and those at high risk of bleeding (4)
- Use in labor and delivery (4)

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 General Dosing Instructions
- 2.2 Administration
- 2.3 Adult Patients < 65 Years of Age
- 2.4 Reduced Doses for Special Populations
- 2.5 Discard Used SPRIX Bottle after 24 Hours

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Cardiovascular Thrombotic Events
- 5.2 Gastrointestinal Bleeding, Ulceration, and Perforation
- 5.3 Hepatotoxicity
- 5.4 Hypertension
- 5.5 Heart Failure and Edema
- 5.6 Renal Toxicity and Hyperkalemia
- 5.7 Anaphylactic Reactions
- 5.8 Exacerbation of Asthma Related to Aspirin Sensitivity
- 5.9 Serious Skin Reactions
- 5.10 Premature Closure of Fetal Ductus Arteriosus
- 5.11 Hematologic Toxicity
- 5.12 Masking of Inflammation and Fever
- 5.13 Laboratory Monitoring

FULL PRESCRIBING INFORMATION

WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS

Cardiovascular Thrombotic Events

- **Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in treatment and may increase with duration of use [see Warnings and Precautions (5.1)].**
- **SPRIX® is contraindicated in the setting of coronary artery bypass graft (CABG) surgery [see Contraindications (4) and Warnings and Precautions (5.1)].**

Gastrointestinal Bleeding, Ulceration, and Perforation

- **NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious GI events [see Warnings and Precautions (5.2)].**

1 INDICATIONS AND USAGE

SPRIX is indicated in adult patients for the short term (up to 5 days) management of moderate to moderately severe pain that requires analgesia at the opioid level.

Limitations of Use

- Sprix is not for use in pediatric patients less than 2 years of age.

2 DOSAGE AND ADMINISTRATION

2.1 General Dosing Instructions

Use the lowest effective dosage for the shortest duration consistent with individual patient treatment goals [see Warnings and Precautions (5.1)].
The total duration of use of SPRIX alone or sequentially with other formulations of ketorolac (IM/IV or oral) must not exceed 5 days because of the potential for increasing the frequency and severity of adverse reactions associated with the recommended doses [see Warnings and Precautions (5.15)].

Do not use SPRIX concomitantly with other formulations of ketorolac or other NSAIDs [see Warnings and Precautions (5.15)].

2.2 Administration

SPRIX is not an inhaled product. Do not inhale when administering this product.

Instruct patients to administer as follows:

1. First hold the finger flange with fingers, and remove the clear plastic cover with opposite hand; then remove the blue plastic safety clip. Keep the clear plastic cover, and throw away the blue plastic safety clip.
2. Before using the bottle for the **FIRST** time, activate the pump. To activate the pump, hold the bottle at arm's length away from the body with index finger and middle finger resting on the top of the finger flange and thumb supporting the base.

Press down evenly and release the pump 5 times. Patient may not see a spray the first few times he/she presses down.

The bottle is now ready to use. There is no need to activate the pump again if more doses are used from the bottle.

3. It's important to get the medication to the correct place in the nose so it will be most effective.

- Blow nose gently to clear nostrils.
- Sit up straight or stand. Tilt head slightly forward.
- Insert the tip of the container into your right nostril.
- Point the container away from the center of your nose.
- Hold your breath and spray once into your right nostril, pressing down evenly on both sides.
- Immediately after administration, resume breathing through mouth to reduce expelling the product. Also pinch the nose to help retain the spray if it starts to drip.

If only one spray per dose is prescribed, administration is complete; skip to Step 5 below.

4. If a dose of 2 sprays is prescribed, repeat the process in Step 3 for the left nostril. Again, be sure to point the spray away from the center of nose. Spray once into the left nostril.

5. Replace the clear plastic cover and place the bottle in a cool, dry location out of direct sunlight, such as inside a medication cabinet. Keep out of reach of children.

2.3 Adult Patients < 65 Years of Age

The recommended dose is 31.5 mg SPRIX (one 15.75 mg spray in each nostril) every 6 to 8 hours. The maximum daily dose is 126 mg (four doses).

2.4 Reduced Doses for Special Populations

For patients ≥ 65 years of age, renally impaired patients, and adult patients less than 50 kg (110 lbs), the recommended dose is 15.75 mg SPRIX (one 15.75 mg spray in **only one** nostril) every 6 to 8 hours. The maximum daily dose is 63 mg (four doses) [see Warnings and Precautions (5.2, 5.6)].

2.5 Discard Used SPRIX Bottle after 24 Hours

Do not use any single SPRIX bottle for more than one day as it will not deliver the intended dose after 24 hours. Therefore, the bottle must be discarded no more than 24 hours after taking the first dose, even if the bottle still contains some liquid.

3 DOSAGE FORMS AND STRENGTHS

SPRIX (ketorolac tromethamine) Nasal Spray: 15.75 mg of ketorolac tromethamine in each 100 µL spray. Each 1.7 g bottle contains 8 sprays.

4 CONTRAINDICATIONS

SPRIX is contraindicated in the following patients:

- Known hypersensitivity (e.g., anaphylactic reactions and serious skin reactions) to ketorolac or any components of the drug product [see Warnings and Precautions (5.7, 5.9)]
- History of asthma, urticaria, or other allergic-type reactions after taking aspirin or other NSAIDs. Severe, sometimes fatal, anaphylactic reactions to NSAIDs have been reported in such patients [see Warnings and Precautions (5.7, 5.8)]
- In the setting of coronary artery bypass graft (CABG) surgery [see Warnings and Precautions (5.1)]

Use in patients with active peptic ulcer disease and in patients with recent gastrointestinal bleeding or perforation [see Warnings and Precautions (5.2)]

Use as a prophylactic analgesic before any major surgery [see Warnings and Precautions (5.11)]

Use in patients with advanced renal disease or patients at risk for renal failure due to volume depletion [see Warnings and Precautions (5.6)]

Use in labor and delivery. Through its prostaglandin synthesis inhibitory effect, ketorolac may adversely affect fetal circulation and inhibit uterine contractions, thus increasing the risk of uterine ischemia [see Use in Specific Populations (8.1)]

Use in patients with suspected or confirmed cerebrovascular bleeding, hemorrhagic diathesis, incomplete hemostasis, or those for whom hemostasis is critical [see Warnings and Precautions (5.11), Drug Interactions (7)]

Concomitant use with probenecid [see Drug Interactions (7)]

Concomitant use with pentoxifylline [see Drug Interactions (7)]

WARNINGS AND PRECAUTIONS

- **Hepatotoxicity:** Inform patients of warning signs and symptoms of hepatotoxicity. Discontinue if abnormal liver tests persist or worsen or if clinical signs and symptoms of liver disease develop. (5.3)
- **Hypertension:** Patients taking some antihypertensive medications may have impaired response to these therapies when taking NSAIDs. Monitor blood pressure. (5.4, 7)
- **Heart Failure and Edema:** Avoid use of SPRIX in patients with severe heart failure unless benefits are expected to outweigh risk of worsening heart failure. (5.5)
- **Renal Toxicity:** Monitor renal function in patients with renal or hepatic impairment, heart failure, dehydration, or hypovolemia. Avoid use of SPRIX in patients with advanced renal disease unless benefits are expected to outweigh risk of worsening renal function. (5.6)
- **Anaphylactic Reactions:** Seek emergency help if an anaphylactic reaction occurs. (5.7)
- **Exacerbation of Asthma Related to Aspirin Sensitivity:** SPRIX is contraindicated in patients with aspirin-sensitive asthma. Monitor patients with preexisting asthma (without aspirin sensitivity). (5.8)
- **Serious Skin Reactions:** Discontinue SPRIX at first appearance of skin rash or other signs of hypersensitivity. (5.9)
- **Premature Closure of Fetal Ductus Arteriosus:** Avoid use in pregnant women starting at 30 weeks gestation. (5.10, 8.1)
- **Hematologic Toxicity:** Monitor hemoglobin or hematocrit in patients with any signs or symptoms of anemia. Do not use SPRIX in patients for whom hemostasis is critical. (5.11, 7)
- **Limitations of Use:** SPRIX should not be used concomitantly with IM/IV or oral ketorolac, aspirin, or other NSAIDs. (5.15)

ADVERSE REACTIONS

Most common adverse reactions (incidence ≥2%) in patients treated with SPRIX and occurring at a rate at least twice that of placebo are nasal discomfort, rhinalgia, increased lacrimation, throat irritation, oliguria, rash, bradycardia, decreased urine output, increased ALT and/or AST, hypertension, and rhinitis. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact **Egalet US Inc. at 1-800-518-1084 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

DRUG INTERACTIONS

- **Drugs that Interfere with Hemostasis (e.g., warfarin, aspirin, SSRIs/SNRIs):** Monitor patients for bleeding who are concomitantly taking SPRIX with drugs that interfere with hemostasis. Concomitant use of SPRIX and analgesic doses of aspirin is not generally recommended. (7)
- **ACE Inhibitors, Angiotensin Receptor Blockers (ARB), or Beta-Blockers:** Concomitant use with SPRIX may diminish the antihypertensive effect of these drugs. Monitor blood pressure. (7)
- **ACE Inhibitors and ARBs:** Concomitant use with SPRIX in elderly, volume depleted, or those with renal impairment may result in deterioration of renal function. In such high risk patients, monitor for signs of worsening renal function. (7)
- **Diuretics:** NSAIDs can reduce natriuretic effect of furosemide and thiazide diuretics. Monitor patients to assure diuretic efficacy including antihypertensive effects. (7)
- **Digoxin:** Concomitant use with SPRIX can increase serum concentration and prolong half-life of digoxin. Monitor serum digoxin levels. (7)

USE IN SPECIFIC POPULATIONS

Pregnancy: Use of NSAIDs during the third trimester of pregnancy increases the risk of premature closure of the fetal ductus arteriosus. Avoid use of NSAIDs in pregnant women starting at 30 weeks gestation. (5.10, 8.1)

Infertility: NSAIDs are associated with reversible infertility. Consider withdrawal of SPRIX in women who have difficulties conceiving. (8.3)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 01/2018

5.14	Eye Exposure
5.15	Limitations of Use
6	ADVERSE REACTIONS
6.1	Clinical Trials Experience
6.2	Postmarketing Experience
7	DRUG INTERACTIONS
8	USE IN SPECIFIC POPULATIONS
8.1	Pregnancy
8.2	Lactation
8.3	Females and Males of Reproductive Potential
8.4	Pediatric Use
8.5	Geriatric Use
10	OVERDOSAGE
11	DESCRIPTION
12	CLINICAL PHARMACOLOGY
12.1	Mechanism of Action
12.3	Pharmacokinetics
13	NONCLINICAL TOXICOLOGY
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
14	CLINICAL STUDIES
14.1	Postoperative Pain
16	HOW SUPPLIED/STORAGE AND HANDLING
17	PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

5 WARNINGS AND PRECAUTIONS

5.1 Cardiovascular Thrombotic Events

Clinical trials of several COX-2 selective and nonselective NSAIDs of up to three years duration have shown an increased risk of serious cardiovascular (CV) thrombotic events, including myocardial infarction (MI) and stroke, which can be fatal. Based on available data, it is unclear that the risk for CV thrombotic events is similar for all NSAIDs. The relative increase in serious CV thrombotic events over baseline conferred by NSAID use appears to be similar in those with and without known CV disease or risk factors for CV disease. However, patients with known CV disease or risk factors had a higher absolute incidence of excess serious CV thrombotic events, due to their increased baseline rate. Some observational studies found that this increased risk of serious CV thrombotic events began as early as the first weeks of treatment. The increase in CV thrombotic risk has been observed most consistently at higher doses.

To minimize the potential risk for an adverse CV event in NSAID-treated patients, use the lowest effective dose for the shortest duration possible. Physicians and patients should remain alert for the development of such events, throughout the entire treatment course, even in the absence of previous CV symptoms. Patients should be informed about the symptoms of serious CV events and the steps to take if they occur. There is no consistent evidence that concurrent use of aspirin mitigates the increased risk of serious CV thrombotic events associated with NSAID use. The concurrent use of aspirin and an NSAID, such as ketorolac, increases the risk of serious gastrointestinal (GI) events [see Warnings and Precautions (5.2)].

Status Post Coronary Artery Bypass Graft (CABG) Surgery

Two large, controlled clinical trials of a COX-2 selective NSAID for the treatment of pain in the first 10–14 days following CABG surgery found an increased incidence of myocardial infarction and stroke. NSAIDs are contraindicated in the setting of CABG [see Contraindications (4)].

Post-MI Patients

Observational studies conducted in the Danish National Registry have demonstrated that patients treated with NSAIDs in the post-MI period were at increased risk of reinfarction, CV-related death, and all-cause mortality beginning in the first week of treatment. In this same cohort, the incidence of death in the first year post-MI was 20 per 100 person years in NSAID-treated patients compared to 12 per 100 person years in non-NSAID exposed patients. Although the absolute rate of death declined somewhat after the first year post-MI, the increased relative risk of death in NSAID users persisted over at least the next four years of follow-up.

Avoid the use of SPRIX in patients with a recent MI unless the benefits are expected to outweigh the risk of recurrent CV thrombotic events. If SPRIX is used in patients with a recent MI, monitor patients for signs of cardiac ischemia.

5.2 Gastrointestinal Bleeding, Ulceration, and Perforation

SPRIX is contraindicated in patients with active peptic ulcers and/or GI bleeding and in patients with recent gastrointestinal bleeding or perforation [see Contraindications (4)]. NSAIDs, including ketorolac, cause serious gastrointestinal (GI) adverse events including inflammation, bleeding, ulceration, and perforation of the esophagus, stomach, small intestine, or large intestine, which can be fatal. These serious adverse events can occur at any time, with or without warning symptoms, in patients treated with NSAIDs. Only one in five patients who develop a serious upper GI adverse event on NSAID therapy is symptomatic. Upper GI ulcers, gross bleeding, or perforation caused by NSAIDs occurred in approximately 1% of patients treated for 3–6 months, and in about 2%–4% of patients treated for one year. However, even short-term NSAID therapy is not without risk.

Risk Factors for GI Bleeding, Ulceration, and Perforation

Patients with a prior history of peptic ulcer disease and/or GI bleeding who used NSAIDs had a greater than 10-fold increased risk for developing a GI bleed compared to patients without these risk factors. Other factors that increase the risk of GI bleeding in patients treated with NSAIDs include longer duration of NSAID therapy; concomitant use of oral corticosteroids, aspirin, anticoagulants, or selective serotonin reuptake inhibitors (SSRIs); smoking; use of alcohol; older age; and poor general health status. Most postmarketing reports of fatal GI events occurred in elderly or debilitated patients. Additionally, patients with advanced liver disease and/or coagulopathy are at increased risk for GI bleeding.

Strategies to Minimize the GI Risks in NSAID-Treated Patients:

- Use the lowest effective dosage for the shortest possible duration.
- Avoid administration of more than one NSAID at a time.
- Avoid use in patients at higher risk unless benefits are expected to outweigh the increased risk of bleeding. For such patients, consider alternate therapies other than NSAIDs. Do not use Sprix in those with active GI bleeding.
- Remain alert for signs and symptoms of GI ulceration and bleeding during NSAID therapy.
- If a serious GI adverse event is suspected, promptly initiate evaluation and treatment, and discontinue SPRIX until a serious GI adverse event is ruled out.
- In the setting of concomitant use of low-dose aspirin for cardiac prophylaxis, monitor patients more closely for evidence of GI bleeding [see Drug Interactions (7)].
- Use great care when giving SPRIX to patients with a history of inflammatory bowel disease (ulcerative colitis, Crohn's disease) as their condition may be exacerbated.

5.3 Hepatotoxicity

Elevations of ALT or AST (three or more times the upper limit of normal [ULN]) have been reported in approximately 1% of NSAID-treated patients in clinical trials. In addition, rare, sometimes fatal, cases of severe hepatic injury, including fulminant hepatitis, liver necrosis, and hepatic failure have been reported. Elevations of ALT or AST (less than three times ULN) may occur in up to 15% of patients treated with NSAIDs including ketorolac.

Inform patients of the warning signs and symptoms of hepatotoxicity (e.g., nausea, fatigue, lethargy, diarrhea, pruritus, jaundice, right upper quadrant tenderness, and "flu-like" symptoms). If clinical signs and symptoms consistent with liver disease develop, or if systemic manifestations occur (e.g., eosinophilia, rash, etc.), discontinue SPRIX immediately, and perform a clinical evaluation of the patient.

5.4 Hypertension

NSAIDs, including SPRIX, can lead to new onset of hypertension or worsening of preexisting hypertension, either of which may contribute to the increased incidence of CV events. Patients taking antihypertensive medication should be closely followed (see Drug Interactions (7)). Monitor blood pressure (BP) during the initiation of NSAID treatment and throughout the course of therapy.

5.5 Heart Failure and Edema

The Coxib and traditional NSAID Trialists' Collaboration meta-analysis of randomized controlled trials demonstrated an approximately two-fold increase in hospitalizations for heart failure in COX-2 selective-treated patients and nonselective NSAID-treated patients compared to placebo-treated patients. In a Danish National Registry study of patients with heart failure, NSAID use increased the risk of MI, hospitalization for heart failure, and death. Additionally, fluid retention and edema have been observed in some patients treated with NSAIDs. Use of ketorolac may blunt the CV effects of several therapeutic agents used to treat these medical conditions (e.g., diuretics, ACE inhibitors, or angiotensin receptor blockers [ARBs]) [see Drug Interactions (7)].

Avoid the use of SPRIX in patients with severe heart failure unless the benefits are expected to outweigh the risk of worsening heart failure. If SPRIX is used in patients with severe heart failure, monitor patients for signs of worsening heart failure.

5.6 Renal Toxicity and Hyperkalemia

Ketorolac and its metabolites are eliminated primarily by the kidneys. Patients with reduced creatinine clearance will have diminished clearance of the drug [see Clinical Pharmacology (12.3)]. SPRIX is contraindicated in patients with advanced renal impairment [see Contraindications (4)].

Renal Toxicity

Long-term administration of NSAIDs has resulted in renal papillary necrosis and other renal injury. Renal toxicity has also been seen in patients in whom renal prostaglandins have a compensatory role in the maintenance of renal perfusion. In these patients, administration of a NSAID may cause a dose-dependent reduction in prostaglandin formation and, secondarily, in renal blood flow, which may precipitate overt renal decompensation. Patients at greatest risk of this reaction are those with impaired renal function, dehydration, hypovolemia, heart failure, liver dysfunction, those taking diuretics and ACE inhibitors or ARBs, and the elderly. Discontinuation of NSAID therapy is usually followed by recovery to the pretreatment state.

No information is available from controlled clinical studies regarding the use of SPRIX in patients with advanced renal disease. The renal effects of SPRIX may hasten the progression of renal dysfunction in patients with preexisting renal disease. Correct volume status in dehydrated or hypovolemic patients prior to initiating SPRIX. Monitor renal function in patients with renal or hepatic impairment, heart failure, dehydration, or hypovolemia during use of SPRIX [see Drug Interactions (7)]. Avoid the use of SPRIX in patients with advanced renal disease unless the benefits are expected to outweigh the risk of worsening renal function. If SPRIX is used in patients with advanced renal disease, monitor patients for signs of worsening renal function.

Hyperkalemia

Increases in serum potassium concentration, including hyperkalemia, have been reported with use of NSAIDs, even in some patients without renal impairment. In patients with normal renal function, these effects have been attributed to a hyporeninemic-hypoaldosteronism state.

5.7 Anaphylactic Reactions

Ketorolac has been associated with anaphylactic reactions in patients with and without known hypersensitivity to ketorolac and in patients with aspirin-sensitive asthma [see Contraindications (4) and Warnings and Precautions (5.8)]. Seek emergency help if an anaphylactic reaction occurs.

5.8 Exacerbation of Asthma Related to Aspirin Sensitivity

A subpopulation of patients with asthma may have aspirin-sensitive asthma which may include chronic rhinosinusitis complicated by nasal polyps, severe, potentially fatal bronchospasm, and/or intolerance to aspirin and other NSAIDs. Because cross-reactivity between aspirin and other NSAIDs has been reported in such aspirin-sensitive patients, SPRIX is contraindicated in patients with this form of aspirin sensitivity [see Contraindications (4)]. When SPRIX is used in patients with preexisting asthma (without known aspirin sensitivity), monitor patients for changes in the signs and symptoms of asthma.

5.9 Serious Skin Reactions

NSAIDs, including ketorolac, can cause serious skin adverse reactions such as exfoliative dermatitis, Stevens-Johnson Syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. These serious events may occur without warning. Inform patients about the signs and symptoms of serious skin reactions, and to discontinue the use of SPRIX at the first appearance of skin rash or any other sign of hypersensitivity. SPRIX is contraindicated in patients with previous serious skin reactions to NSAIDs [see Contraindications (4)].

5.10 Premature Closure of Fetal Ductus Arteriosus

Ketorolac may cause premature closure of the fetal ductus arteriosus. Avoid use of NSAIDs, including SPRIX, in pregnant women starting at 30 weeks of gestation (third trimester) [see Use in Specific Populations (8.1)].

5.11 Hematologic Toxicity

Anemia has occurred in NSAID-treated patients. This may be due to occult or gross blood loss, fluid retention, or an incompletely described effect upon erythropoiesis. If a patient treated with SPRIX has any signs or symptoms of anemia, monitor hemoglobin or hematocrit. Do not use SPRIX in patients for whom hemostasis is critical [see Contraindications (4), Drug Interactions (7)].

NSAIDs, including SPRIX, may increase the risk of bleeding events. Co-morbid conditions such as coagulation disorders or concomitant use of warfarin, other anticoagulants, antiplatelet agents (e.g., aspirin), serotonin reuptake inhibitors (SSRIs) and serotonin norepinephrine reuptake inhibitors (SNRIs) may increase this risk. Monitor these patients for signs of bleeding [see Drug Interactions (7)].

The concurrent use of ketorolac and therapy that affects hemostasis, including prophylactic low dose heparin (2500 to 5000 units q12h), warfarin and dextrans, has not been studied extensively, but may also be associated with an increased risk of bleeding. Until data from such studies are available, carefully weigh the benefits against the risks and use such concomitant therapy only with extreme caution. Monitor patients receiving therapy that affects hemostasis closely.

In clinical trials, serious adverse events related to bleeding were more common in patients treated with SPRIX than placebo. In clinical trials and in postmarketing experience with ketorolac IV and IM dosing, postoperative hematomas and other signs of wound bleeding have been reported in association with peri-operative use. Therefore, use SPRIX with caution in the postoperative setting when hemostasis is critical.

5.12 Masking of Inflammation and Fever

The pharmacological activity of SPRIX in reducing inflammation, and possibly fever, may diminish the utility of diagnostic signs in detecting infections.

5.13 Laboratory Monitoring

Because serious GI bleeding, hepatotoxicity, and renal injury can occur without warning symptoms or signs, consider monitoring patients on long-term NSAID treatment with a CBC and a chemistry profile periodically [see Warnings and Precautions (5.2, 5.3, 5.6)].

5.14 Eye Exposure

Avoid contact of SPRIX with the eyes. If eye contact occurs, wash out the eye with water or saline, and consult a physician if irritation persists for more than an hour.

5.15 Limitations of Use

The total duration of use of SPRIX alone or sequentially with other forms of ketorolac is not to exceed 5 days. SPRIX must not be used concomitantly with other forms of ketorolac or other NSAIDs [see Dosage and Administration (2.1)].

6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- Cardiovascular Thrombotic Events [see Warnings and Precautions (5.1)]
- GI Bleeding, Ulceration and Perforation [see Warnings and Precautions (5.2)]
- Hepatotoxicity [see Warnings and Precautions (5.3)]
- Hypertension [see Warnings and Precautions (5.4)]
- Heart Failure and Edema [see Warnings and Precautions (5.5)]
- Renal Toxicity and Hyperkalemia [see Warnings and Precautions (5.6)]
- Anaphylactic Reactions [see Warnings and Precautions (5.7)]
- Serious Skin Reactions [see Warnings and

treated with NSAIDs has also shown a reversible delay in ovulation. Consider withdrawal of NSAIDs, including SPRIX, in women who have difficulties conceiving or who are undergoing investigation of infertility.

8.4 Pediatric Use

SPRIX is not for use in pediatric patients less than 2 years of age. The safety and effectiveness of ketorolac in pediatric patients 17 years of age and younger have not been established.

8.5 Geriatric Use

Exercise caution when treating the elderly (65 years and older) with SPRIX. Elderly patients, compared to younger patients, are at greater risk for NSAID-associated serious cardiovascular, gastrointestinal, and/or renal adverse reactions. If the anticipated benefit for the elderly patient outweighs these potential risks, start dosing at the low end of the dosing range, and monitor patients for adverse effects [see *Dosage and Administration* (2.4), *Warnings and Precautions* (5.1, 5.2, 5.3, 5.6, 5.13), *Clinical Pharmacology* (12.3)]. After observing the response to initial therapy with SPRIX, adjust the dose and frequency to suit an individual patient's needs.

Ketorolac and its metabolites are known to be substantially excreted by the kidneys, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, use caution in this patient population, and it may be useful to monitor renal function [see *Clinical Pharmacology* (12.3)].

10 OVERDOSAGE

Symptoms following acute NSAID overdosages have been typically limited to lethargy, drowsiness, nausea, vomiting, and epigastric pain, which have been generally reversible with supportive care. Gastrointestinal bleeding has occurred. Hypertension, acute renal failure, respiratory depression, and coma have occurred, but were rare [see *Warnings and Precautions* (5.1, 5.2, 5.4, 5.6)].


There has been no experience with overdosage of SPRIX. In controlled overdosage studies with IM ketorolac injection, daily doses of 300 mg given for five days (approximately 9 times the maximum daily dose of SPRIX) caused abdominal pain and peptic ulcers, which healed after discontinuation of dosing. Single overdoses of ketorolac tromethamine have been variably associated with abdominal pain, nausea, vomiting, hyperventilation, peptic ulcers and/or erosive gastritis, and renal dysfunction.

Manage patients with symptomatic and supportive care following an NSAID overdosage. There are no specific antidotes. Consider emesis and/or activated charcoal (60 to 100 grams in adults, 1 to 2 grams per kg of body weight in pediatric patients) and/or osmotic cathartic in symptomatic patients seen within four hours of ingestion or in patients with a large overdosage (5 to 10 times the recommended dosage). Forced diuresis, alkalization of urine, hemodialysis, or hemoperfusion may not be useful due to high protein binding. For additional information about overdosage treatment contact a poison control center (1-800-222-1222).

11 DESCRIPTION

SPRIX (ketorolac tromethamine) Nasal Spray is a member of the pyrrolo-pyrrole group of nonsteroidal anti-inflammatory drugs, available as a clear, colorless to yellow solution packaged in a glass vial with a snap on spray pump that delivers 15.75 mg ketorolac tromethamine per spray and is intended for intranasal administration. The chemical name is (+)-5-benzoyl-2,3-dihydro-1H-pyrrolo[1,2-c]pyridine-1-carboxylic acid, compound with 2-amino-2-(hydroxymethyl)-1,3-propanediol (1:1). The molecular weight is 376.41.

Its molecular formula is C₁₇H₁₉N₃O₃(C₉H₉NO₃•C₈H₁₁NO₃), and it has the following chemical structure.



The inactive ingredients in SPRIX include: edetate disodium (EDTA), monobasic potassium phosphate, sodium hydroxide, and water for injection.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Ketorolac has analgesic, anti-inflammatory, and antipyretic properties. The mechanism of action of SPRIX, like that of other NSAIDs, is not completely understood but involves inhibition of cyclooxygenase (COX-1 and COX-2), an early component of the arachidonic acid cascade, resulting in the reduced synthesis of prostaglandins, thromboxanes, and prostacyclin.

Ketorolac is a potent inhibitor of prostaglandin synthesis *in vitro*. Ketorolac concentrations reached during therapy have produced *in vivo* effects. Prostaglandins sensitize afferent nerves and potentiate the action of bradykinin in inducing pain in animal models. Prostaglandins are mediators of inflammation. Because ketorolac is an inhibitor of prostaglandin synthesis, its mode of action may be due to a decrease of prostaglandins in peripheral tissues.

12.3 Pharmacokinetics

The half-lives of ketorolac by the IN and IM routes were similar. The bioavailability of ketorolac by the IN route of administration of a 31.5 mg dose was approximately 60% compared to IM administration. (See Table 3).

Table 3: Pharmacokinetic Parameters of Ketorolac Tromethamine after Intramuscular (IM) and Intranasal (IN) Administration

Ketorolac Tromethamine	C _{max} (SD) ng/mL	t _{max} (range) hours	AUC _{0-∞} (SD) ng•h/mL	T _{1/2} (SD) hours
30 mg IM	2382.2 (432.7)	0.75 (0.25-1.03)	11152.8 (4260.1)	4.80 (1.18)
31.5 mg IN (SPRIX)	1805.8 (882.8)	0.75 (0.50-2.00)	7477.3 (3654.4)	5.24 (1.33)
15 mg IM	1163.4 (279.9)	0.75 (0.25-1.50)	5196.3 (2076.7)	5.00 (1.72)

C_{max} = maximum plasma concentration; t_{max} = time of C_{max}; AUC_{0-∞} = complete area under the concentration-time curve; T_{1/2} = half-life; SD = standard deviation. All values are means, except t_{max}, for which medians are reported.

Absorption

In a study in which SPRIX (31.5 mg) was administered to healthy volunteers four times daily for 5 days, the C_{max}, t_{max}, and AUC values following the final dose were comparable to those obtained in the single-dose study. Accumulation of ketorolac has not been studied in special populations, geriatric, pediatric, renal failure or hepatic disease patients.

Distribution

Scintigraphic assessment of drug disposition of ketorolac following SPRIX intranasal dosing demonstrated that most of the ketorolac was deposited in the nasal cavity and pharynx, with less than 20% deposited in the esophagus and stomach, and zero or negligible deposition in the lungs (<0.5%).

The mean apparent volume (V_D) of ketorolac tromethamine following complete distribution was approximately 13 liters. This parameter was determined from single-dose data. The ketorolac tromethamine racemate has been shown to be highly protein bound (99.2%). Nevertheless, plasma concentrations as high as 10 mcg/mL will only occupy approximately 5% of the albumin binding sites. Thus, the unbound fraction for each enantiomer will be constant over the therapeutic range. A decrease in serum albumin, however, will result in increased free drug concentrations. Therapeutic concentrations of dioxin, warfarin, ibuprofen, naproxen, piroxicam, acetaminophen, phenytoin, and tolbutamide did not alter ketorolac protein binding. *In vitro* studies indicate that, at therapeutic concentrations of salicylate (300 mcg/mL), the binding of ketorolac was reduced from approximately 99.2% to 97.5%, representing a potential twofold increase in unbound ketorolac plasma levels. The *in vitro* binding of warfarin to plasma proteins is only slightly reduced by ketorolac (99.5% control vs. 99.3%) when ketorolac plasma concentrations reach 5 to 10 mcg/mL. Ketorolac tromethamine is excreted in human milk.

Elimination

Metabolism

Ketorolac tromethamine is largely metabolized in the liver. The metabolic products are hydroxylated and conjugated forms of the parent drug. The products of metabolism, and some unchanged drug, are excreted in the urine. There is no evidence in animal or human studies that ketorolac induces or inhibits hepatic enzymes capable of metabolizing itself or other drugs.

Excretion

The principal route of elimination of ketorolac and its metabolites is renal. About 92% of a given dose is found in the urine, approximately 40% as metabolites and 60% as unchanged ketorolac. Approximately 6% of a dose is excreted in the feces. A single-dose study with 10 mg ketorolac tromethamine (n = 9) demonstrated that the S-enantiomer is cleared approximately two times faster than the R-enantiomer and that the clearance was independent of the route of administration. This means that the ratio of S/R plasma concentrations decreases with time after each dose. There is little or no inversion of the R- to S- form in humans.

The half-life of the ketorolac tromethamine S-enantiomer was approximately 2.5 hours (SD ± 0.4) compared with 5 hours (SD ± 1.7) for the R-enantiomer. In other studies, the half-life for the racemate has been reported to lie within the range of 5 to 6 hours.

Specific Populations

Geriatric: A single-dose study was conducted to compare the pharmacokinetics of SPRIX (31.5 mg) in subjects ≥ age 65 to the pharmacokinetics in subjects < age 65. Exposure to ketorolac was increased by 23% for the ≥ 65 population as compared to subjects < 65. Peak concentrations of 2928 and 1840 ng/mL were observed for the elderly and nonelderly adult populations, respectively, at 0.75 h after dosing. In the elderly population a longer terminal half-life was observed as compared to the nonelderly adults (4.5 h vs. 3.3 h, respectively).

Race: Pharmacokinetic differences due to race have not been identified.

Hepatic Impairment: There was no significant difference in estimates of half-life, AUC_{0-∞} and C_{max} in 7 patients with liver disease compared to healthy volunteers.

Renal Impairment: Based on single-dose data only, the mean half-life of ketorolac tromethamine in renally impaired patients is between 6 and 19 hours, and is dependent on the extent of the impairment. There is poor correlation between creatinine clearance and total ketorolac tromethamine clearance in the elderly and populations with renal impairment (r = 0.5).

In patients with renal disease, the AUC_{0-∞} of each enantiomer increased by approximately 100% compared with healthy volunteers. The volume of distribution doubles for the S-enantiomer and increases by 1/5th for the R-enantiomer. The increase in volume of distribution of ketorolac tromethamine implies an increase in unbound fraction. The AUC_{0-∞} ratio of the ketorolac tromethamine enantiomers in healthy subjects and patients remained similar, indicating there was no selective excretion of either enantiomer in patients compared to healthy subjects.

Allergic Rhinitis: Comparison of the pharmacokinetics of SPRIX in subjects with allergic rhinitis to data from a previous study in healthy subjects showed no differences that would be of clinical consequence for the efficacy or safety of SPRIX.

Drug Interaction Studies

Aspirin: When NSAIDs were administered with aspirin, the protein binding of NSAIDs were reduced; although the clearance of free NSAID was not altered. The clinical significance of this interaction is not known. See Table 2 for clinically significant drug interactions of NSAIDs with aspirin [see *Drug Interactions* (7)].

Other Nasal Spray Products: A study was conducted in subjects with symptomatic allergic rhinitis to assess the effects of the commonly used nasal spray products oxymetazoline hydrochloride and fluticasone propionate on the pharmacokinetics of SPRIX. Subjects received a single dose of oxymetazoline nasal spray followed by a single dose (31.5 mg) of SPRIX 30 min later. Subjects also received fluticasone nasal spray (200 mcg as 2 x 50 mcg in each nostril) for seven days, with a single dose (31.5 mg) of SPRIX on the 7th day. Administration of these common intranasal products had no effect of clinical significance on the rate or extent of ketorolac absorption.

Probenecid: Concomitant administration of oral ketorolac and probenecid resulted in decreased clearance and volume of distribution of ketorolac and significant increases in ketorolac plasma levels (total AUC increased approximately threefold from 5.4 to 17.8 mcg•h/mL), and terminal half-life increased approximately twofold from 6.6 to 15.1 hours [see *Drug Interactions* (7)].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

An 18-month study in mice with oral doses of ketorolac at 2 mg/kg/day (approximately 1/3 times the human systemic exposure at the recommended maximum IN dose of 31.5 mg four times a day, based on area-under-the-plasma-concentration curve [AUC]), and a 24-month study in rats at 5 mg/kg/day (approximately 0.8 times the human AUC) showed no evidence of tumorigenicity.

Mutagenesis

Ketorolac was not mutagenic in the Ames test, unscheduled DNA synthesis and repair, or in forward mutation assays. Ketorolac did not cause chromosome breakage in the *in vivo* mouse micronucleus assay. At 1590 µg/mL and at higher concentrations, ketorolac increased the incidence of chromosomal aberrations in Chinese hamster ovarian cells.

Impairment of fertility

Impairment of fertility did not occur in male or female rats at oral doses of 9 mg/kg (approximately 1.3 times the human AUC) and 16 mg/kg (approximately 2.4 times the human AUC) of ketorolac, respectively.

14 CLINICAL STUDIES

14.1 Postoperative Pain

The effect of SPRIX on acute pain was evaluated in two multi-center, randomized, double-blind, placebo-controlled studies.

In a study of adults who had undergone elective abdominal or orthopedic surgery, 300 patients were randomized and treated with SPRIX or placebo administered every 6 hours and morphine administered via patient controlled analgesia on an as needed basis. Efficacy was demonstrated as a statistically significant greater reduction in the summed pain intensity difference over 48 hours in patients who received SPRIX as compared to those receiving placebo. The clinical relevance of this is reflected in the finding that patients treated with SPRIX required 36% less morphine over 48 hours than patients treated with placebo.

In a study of adults who had undergone elective abdominal surgery, 321 patients were randomized and treated with SPRIX or placebo administered every 6 hours and morphine administered via patient controlled analgesia on an as needed basis. Efficacy was demonstrated as a statistically significant greater reduction in the summed pain intensity difference over 48 hours in patients who received SPRIX as compared to those receiving placebo. The clinical relevance of this is reflected in the finding that patients treated with SPRIX required 26% less morphine over 48 hours than patients treated with placebo.

16 HOW SUPPLIED/STORAGE AND HANDLING

SPRIX (ketorolac tromethamine) Nasal Spray, 15.75 mg/spray, are single-day preservative-free spray bottles, supplied as:

NDC 69344-144-43 Carton containing 5 single-day nasal spray bottles

NDC 69344-144-53 Carton containing 1 single-day nasal spray bottle

Each single-day nasal spray bottle contains a sufficient quantity of solution to deliver 8 sprays for a total of 126 mg of ketorolac tromethamine. Each spray delivers 15.75 mg of ketorolac tromethamine. The delivery system is designed to administer precisely metered doses of 100 µL per spray.

Storage

Protect from light and freezing. Store unopened SPRIX between 2°C to 8°C (36°F to 46°F). During use, keep containers of SPRIX Nasal Spray at controlled room temperature, between 15°C to 30°C (59°F to 86°F), out of direct sunlight. Bottles of SPRIX should be discarded within 24 hours of priming.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide and Instructions for Use) that accompanies each prescription dispensed. Instruct all patients to read and closely follow the FDA-approved SPRIX Patient Instructions to ensure proper administration of SPRIX. When prescribing SPRIX, inform patients or their caregivers of the potential risks of ketorolac treatment, instruct patients to seek medical advice if they develop treatment-related adverse events, advise patients not to give SPRIX to other family members, and advise patients to discard any unused drug. Inform patients, families, or their caregivers of the following information before initiating therapy with SPRIX and periodically during the course of ongoing therapy.

Cardiovascular Thrombotic Events

Advise patients to be alert for the symptoms of cardiovascular thrombotic events, including chest pain, shortness of breath, weakness, or slurring of speech, and to report any of these symptoms to their health care provider immediately [see *Warnings and Precautions* (5.1)].

Gastrointestinal Bleeding, Ulceration, and Perforation

Advise patients to report symptoms of ulcerations and bleeding, including epigastric pain, dyspepsia, melena, and hematemesis to their health care provider. In the setting of concomitant use of low-dose aspirin for cardiac prophylaxis, inform patients of the increased risk for and the signs and symptoms of GI bleeding [see *Contraindications* (4), *Warnings and Precautions* (5.2)].

Hepatotoxicity

Inform patients of the warning signs and symptoms of hepatotoxicity (e.g., nausea, fatigue, lethargy, pruritus, diarrhea, jaundice, right upper quadrant tenderness, and "flu-like" symptoms). If these occur, instruct patients to seek immediate emergency help if such symptoms occur [see *Warnings and Precautions* (5.3)].

Heart Failure and Edema

Advise patients to be alert for the symptoms of congestive heart failure including shortness of breath, unexplained weight gain, or edema and to seek their healthcare provider if such symptoms occur [see *Warnings and Precautions* (5.5)].

Anaphylactic Reactions

Inform patients of the signs of an anaphylactic reaction (e.g., difficulty breathing, swelling of the face or throat). Instruct patients to seek immediate emergency help if these occur [see *Contraindications* (4) and *Warnings and Precautions* (5.7)].

Serious Skin Reactions

Advise patients to stop SPRIX immediately if they develop any type of rash and to contact their healthcare provider as soon as possible [see *Warnings and Precautions* (5.9)].

Female Fertility

Advise females of reproductive potential who desire pregnancy that NSAIDs, including SPRIX, may be associated with a reversible delay in ovulation [see *Use in Specific Populations* (8.9)].

Fetal Toxicity

Inform pregnant women to avoid use of SPRIX and other NSAIDs starting at 30 weeks gestation because of the risk of the premature closure of the fetal ductus arteriosus [see *Warnings and Precautions* (5.10) and *Use in Specific Populations* (8.7)].

Avoid Concomitant Use of NSAIDs

Inform patients that the concomitant use of SPRIX with other NSAIDs or salicylates (e.g., diflunisal, salsalate) is not recommended due to the increased risk of gastrointestinal toxicity, and little or no increase in efficacy [see *Warnings and Precautions* (5.2) and *Drug Interactions* (7)]. Alert patients that NSAIDs may be present in "over the counter" medications for treatment of colds, fever, or insomnia.

Use of NSAIDs and Low-Dose Aspirin

Inform patients not to use low-dose aspirin concomitantly with SPRIX until they talk to their healthcare provider [see *Drug Interactions* (7)].

Renal Effects

SPRIX is eliminated by the kidneys. Advise patients to maintain adequate fluid intake and request medical advice if urine output decreases significantly [see *Contraindications* (4), *Warnings and Precautions* (5.6)].

Limitations of Use

Instruct patients not to use SPRIX for more than 5 days. Use of SPRIX alone or in combination with any other ketorolac product for more than 5 days increases the risk for serious complications including GI bleeding and renal injury [see *Dosage and Administration* (2)].

Single-Day Container

Instruct patients not to use any single bottle of SPRIX for more than one day [see *Dosage and Administration* (2.5)].

Nasal Discomfort

Advise patients that they may experience transient, mild to moderate nasal irritation or discomfort upon dosing.

Manufactured for and Distributed by:

Egalet US Inc.
Wayne, PA 19087

Issued: Jan/2018

LBL # 101.02

Medication Guide for Nonsteroidal Anti-inflammatory Drugs (NSAIDs)

What is the most important information I should know about medicines called Nonsteroidal Anti-inflammatory Drugs (NSAIDs)?

NSAIDs can cause serious side effects, including:

- Increased risk of a heart attack or stroke that can lead to death. This risk may happen early in treatment and may increase:
 - with increasing doses of NSAIDs
 - with longer use of NSAIDs

Do not take NSAIDs right before or after a heart surgery called a "coronary artery bypass graft (CABG)."

Avoid taking NSAIDs after a recent heart attack, unless your healthcare provider tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.

- Increased risk of bleeding, ulcers, and tears (perforation) of the esophagus (tube leading from the mouth to the stomach), stomach and intestines:
 - anytime during use
 - without warning symptoms
 - that may cause death

The risk of getting an ulcer or bleeding increases with:

- past history of stomach ulcers, or stomach or intestinal bleeding with use of NSAIDs
- taking medicines called "corticosteroids", "anticoagulants", "SSRIs", or "SNRIs"
- increasing doses of NSAIDs
- older age
- longer use of NSAIDs
- poor health
- smoking
- advanced liver disease
- drinking alcohol
- bleeding problems

NSAIDs should only be used:

- exactly as prescribed
- at the lowest dose possible for your treatment
- for the shortest time needed

What are NSAIDs?

NSAIDs are used to treat pain and redness, swelling, and heat (inflammation) from medical conditions such as different types of arthritis, menstrual cramps, and other types of short-term pain.

Who should not take NSAIDs?

Do not take NSAIDs:

- if you have had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAIDs.
- right before or after heart bypass surgery.

Before taking NSAIDs, tell your healthcare provider about all of your medical conditions, including if you:

- have liver or kidney problems
- have high blood pressure
- have asthma
- are pregnant or plan to become pregnant. Talk to your healthcare provider if you are considering taking NSAIDs during pregnancy. **You should not take NSAIDs after 29 weeks of pregnancy.**
- are breastfeeding or plan to breast feed.

Tell your healthcare provider about all of the medicines you take, including prescription or over-the-counter medicines, vitamins or herbal supplements. NSAIDs and some other medicines can interact with each other and cause serious side effects. Do not start taking any new medicine without talking to your healthcare provider first.

What are the possible side effects of NSAIDs?

NSAIDs can cause serious side effects, including:

See "What is the most important information I should know about medicines called Nonsteroidal Anti-inflammatory Drugs (NSAIDs)?"

- new or worse high blood pressure
- heart failure
- liver problems including liver failure
- kidney problems including kidney failure
- low red blood cells (anemia)
- life-threatening skin reactions

- life-threatening allergic reactions
- Other side effects of NSAIDs include:** stomach pain, constipation, diarrhea, gas, heartburn, nausea, vomiting, and dizziness.

Get emergency help right away if you get any of the following symptoms:

- shortness of breath or trouble breathing
- chest pain
- weakness in one part or side of your body
- slurred speech
- swelling of the face or throat

Stop taking your NSAID and call your healthcare provider right away if you get any of the following symptoms:

- nausea
- more tired or weaker than usual
- diarrhea
- itching
- your skin or eyes look yellow
- indigestion or stomach pain
- flu-like symptoms
- vomit blood
- there is blood in your bowel movement or it is black and sticky like tar
- unusual weight gain
- skin rash or blisters with fever
- swelling of the arms, hands and feet

If you take too much of your NSAID, call your healthcare provider or get medical help right away.

These are not all the possible side effects of NSAIDs. For more information, ask your healthcare provider or pharmacist about NSAIDs.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Other information about NSAIDs

- Aspirin is an NSAID but it does not increase the chance of a heart attack. Aspirin can cause bleeding in the brain, stomach, and intestines. Aspirin can also cause ulcers in the stomach and intestines.
- Some NSAIDs are sold in lower doses without a prescription (over-the-counter). Talk to your healthcare provider before using over-the-counter NSAIDs for more than 10 days.

General information about the safe and effective use of NSAIDs

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use NSAIDs for a condition for which it was not prescribed. Do not give NSAIDs to other people, even if they have the same symptoms that you have. It may harm them.

If you would like more information about NSAIDs, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about NSAIDs that is written for health professionals.

Manufactured for: Egalet US Inc., Wayne, PA 19087

Distributed by: Egalet US Inc., Wayne, PA 19087

For more information, go to www.sprix.com or call 1-800-518-1084.

This Medication Guide has been approved by the U.S. Food and Drug Administration. Issued or Revised: May/2016

Instructions for Use

SPRIX® (spriks)
(ketorolac tromethamine)
Nasal Spray

Read this Instructions for Use before you start using SPRIX and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

Important information:

- SPRIX is for use in your nose only. Do not breathe in (inhale) SPRIX.**
- Each SPRIX bottle has enough pain medicine for 1 day.
- Throw away each SPRIX bottle within 24 hours of taking your first dose, even if the bottle still contains unused medicine.

Your healthcare provider has prescribed SPRIX to treat moderate to severe pain.

- Use SPRIX exactly as your healthcare provider tells you to use it.
- Your healthcare provider will tell you how many sprays you should use each time you use SPRIX.
- Do not use SPRIX for more than 5 days. If you still have pain after 5 days, contact your healthcare provider.
- Do not use SPRIX more than every 6 hours.
- It is important that you drink plenty of fluids while you are using SPRIX. Tell your healthcare provider if you urinate less while using SPRIX.

You may have discomfort or irritation in your nose when using SPRIX. This usually lasts for a short time. Do not breathe in (inhale) SPRIX while spraying.

Using SPRIX Nasal Spray

Parts of your SPRIX bottle

